Journée annuelle ARPEM 2022

Fatigue & Sport

Cas cliniques

H 49ans

Fatigue depuis 4 sem

Impression de baisse de performance avec augmentation de la difficulté pour un même effort

F 35ans

Fatigue globale depuis 3 mois

Chronos moins bons en compétition malgré intensification de l'entraînement

Plus souvent malade et irritable

F 20ans

Fatigue et troubles du sommeil depuis peu

Difficultés à tenir le rythme d'entraînement pour un objectif de marathon

Surcompensation

FOR

Functional Overreaching

Image – Soligard T, et al. BJSM 2016

Surentraining

Image – Soligard T, et al. BJSM 2016

Définition

Meeusen R, et al. Med Sci Sports Exerc 2013

Symptômes

Table 2. Symptoms of overtraining syndrome.

Parasympathetic Alterations ^a	Sympathetic Alterations ^b	Other
Fatigue	Insomnia	Anorexia
Depression	Irritability	Weight loss
Bradycardia	Agitation	Lack of mental concentration
Loss of motivation	Tachycardia	Heavy, sore, stiff muscles
	Hypertension	Anxiety
	Restlessness	Awakening unrefreshed

^aMore common in aerobic sports.

^bMore common in anaerobic sports.

Kreher JB, et al. Sports Health 2012

Déséquilibre

Image – Soligard T, et al. BJSM 2016

Charge

Surentrainement fonctionnel

Surentrainement non-fonctionnel

Syndrome de surentrainement

BORG 6-20 original	BORG 1-10 modifié	% FC maximale	Perception	Activité
6	0	50-60%	repos	repos
7			très très facile	
8				
9	1	60-70%	très très facile	marche
10	2		facile	léger jogging
11		3	modéré	jogging
12	4		70-80%	un peu dur
13		5	80-90%	dur
14	6		90-95%	très dur
15		7		
16	8		maximal	
17		9		
18	10			
19				
20				

Récupérer de quoi? La fatigue

Surentraining fonctionnel

Surentraining non-fonctionnel

Syndrome de surentraining

Réhydrater

Recharger/Restocker

Régénérer/Réparer

Ressourcer/Relaxer

Définition

Meeusen R, et al. Med Sci Sports Exerc 2013

Triggers

Table 4. Potential triggers of overtraining syndrome.³⁰

Increased training load without adequate recovery
Monotony of training
Excessive number of competitions
Sleep disturbances
Stressors including personal life (family, relationships) and occupational
Previous illness
Altitude exposure
Heat injury episode
Severe “bonk”

Kreher JB, et al. Sports Health 2012

Cas cliniques

H 49ans

Charge:

- Course à pied 3x/sem
- Cyclisme 1-2x/sem
- Renforcement 1x/sem

Récupération:

- Pas de régime particulier
- Poids stable
- Sommeil bon

Triggers:

- Marié, 1 fils de 4ans
- Bureautique
- Cours d'aérobic 2x/sem et de danse chorégraphique

F 35ans

Charge:

- Athlétisme (demi-fond) 5x/sem, 60-120'
- ↑ intensité depuis 4sem

Récupération:

- Pas de régime particulier
- Poids stable
- Sommeil 6-7h/nuit

Triggers:

- Mariée, 2 enfants
- Entrepreneurse
- Stress au travail

F 20ans

Charge:

- Course à pied 5x/sem
- Natation 1x/sem
- Vélo pour déplacements

Récupération:

- Végétarienne mais fait attention à ses apports
- Réveils nocturnes, dort peu

Triggers:

- Etudiante EPFZ
- Période d'examens

Epidémiologie

Prévalence de 15-60% (OTS)

- 30% chez les athlètes d'endurance non-élites (NFOR/OTS)
- 60% chez les athlètes d'endurance élités (NFOR/OTS)
- Athlètes suisses: 9% de NFOR, 21% de OTS

Le plus souvent dans les sports:

- D'endurance (course à pied, cyclisme, natation)
- A demande physique moyenne à importante

Pas de différence:

- Sports individuels et d'équipe
- Femmes vs Hommes

Figure 1: Accompanying symptoms of overreaching/overtraining for the NOR/FOR and NFOR/OTS groups. *Significant difference between groups ($p < .05$). **Significant difference between groups ($p < .01$).

Cadegiani FA, et al. *BMJ Open Sport Exerc Med* 2019

Cardoos N, et al. *Curr Sports Med Rep* 2015

Meeusen R, et al. *Med Sci Sports Exerc* 2013

Birrer D, et al. *Schweizerische Zeitschrift für Sportmedizin und Sporttraumatologie* 2013

Physiopathologie – Hypothèses

Glycogène/Dysbalance énergétique

Fatigue centrale/BCAA

Glutamine

Stress oxydatif

Système nerveux autonome

Axe hypothalamo-hypophyso-surrénalien

Cytokines/Inflammation

Dysfonction immunitaire

Hypothesis	Theory	Strengths	Weaknesses
Glycogen hypothesis	Decreased glycogen causes fatigue and decreased performance	Low glycogen can be correlated with decreased performance and exercise-induced fatigue	No proven correlation in the literature between low glycogen and overtrained athletes Athletes with normal glycogen levels still become overtrained Does not account for all symptoms
Central fatigue hypothesis	Increased tryptophan uptake in the brain leads to increased 5-HT centrally and mood symptoms	Exercise correlated with increased tryptophan, 5-HT, and fatigue Rats undergoing intense training have increased 5-HT Selective serotonin reuptake inhibitors decrease performance Athletes getting branched chain amino acids supplement had less fatigue	Few studies measure 5-HT directly Mood changes/fatigue subjective and difficult to study Mood/fatigue influenced by many other factors Does not account for all symptoms
Glutamine hypothesis	Decreased glutamine causes immune dysfunction and increased susceptibility to infection	Glutamine does decrease after prolonged exercise In vitro immune cell function is compromised with decreased glutamine Athletes are more susceptible to upper respiratory tract infections after "intense" exercise	In vivo, decreased plasma glutamine not necessarily correlated with decreased bioavailable glutamine Glutamine supplementation does not improve postexercise impairment of immune cells Some studies show low glutamine in athletes with upper respiratory tract infections, and some do not Glutamine can be influenced by many other factors Increased upper respiratory tract infections are seen in most athletes after intense exercise, not just overtrained ones Does not account for all symptoms
Oxidative stress hypothesis	Excessive oxidative stress causes muscle damage and fatigue	Resting markers of oxidative stress are higher in overtrained athletes and increase with exercise Citrate synthase (marker of oxidative capacity) decreases in overreached rats, so they are more susceptible to oxidative stress	Studies have been small Lack of clinically relevant research Does not account for all symptoms
Autonomic nervous system hypothesis	Parasympathetic predominance causes many symptoms of overtraining syndrome	A study showed variability in autonomic nervous system forces (through heart rate variability) with exercise versus rest Decreased heart rate variability with awakening in overtrained athletes suggests disruption of autonomic nervous system modulation	Decreased nocturnal catecholamines in overtrained athletes in some studies; no change or increased in others Studies looking at catecholamine excretion with methodological differences are hard to compare No difference in heart rate variability/autonomic nervous system influence between overtrained and control athletes during sleep, when free of external influences Does not account for all symptoms

Kreher JB, et al. Sports Health 2012

Hypothesis	Theory	Strengths	Weaknesses
Hypothalamic hypothesis	Dysregulation of the hypothalamus and hormonal axes cause many symptoms of overtraining syndrome	Endurance athletes have activation of the hypothalamic-pituitary-adrenal axis compared with controls	Contradictory data in terms of activation of hypothalamic-pituitary-adrenal/hypothalamic-pituitary-gonadal axes in overtrained athletes and levels of ACTH, cortisol, testosterone Other factors can influence hypothalamic-pituitary-adrenal/hypothalamic-pituitary-gonadal axis activation Does not account for all symptoms
Cytokine hypothesis	Inflammation and cytokine release causes most of the above effects and symptoms of overtraining syndrome	Unified theory accounting for many symptoms of overtraining syndrome and “why” it develops Cytokines may act on hypothalamic centers to regulate “sickness” behavior, causing mood changes and fatigue Subacute muscle injury and cytokines decreases glucose transport into muscles, decreases glycogen, and causes fatigue Tryptophan is used to synthesize inflammatory proteins and decreases with systemic inflammation Decreased tryptophan is associated with depressive symptoms Increased cytokine levels are found in depressed patients Giving participants cytokines caused depressive symptoms Cytokines activate the hypothalamic-pituitary-adrenal system (increasing cortisol) and inhibit the hypothalamic-pituitary-gonadal system (decreasing testosterone) Inflammation causes activation of glucose/protein metabolism and decreased glutamine Increased cytokines that favor TH2 lymphocyte activation lead to increased humoral/decreased cell-mediated immunity and more upper respiratory tract infection	Little evidence actually verifying increased cytokines in overtrained athletes No studies looked at long-term responses to training/overtraining One study showed no change in cytokine levels in overtrained cyclists Cytokine studies to date look predominantly at very fit athletes with questionable application to the general population

Kreher JB, et al. Sports Health 2012

Diagnostic

1. Perte de performance sport-spécifique malgré un repos adéquat de 14-21 jours
2. Des signes et symptômes physiques +/- psychologiques compatibles
3. Absence d'autre cause → Diagnostic d'exclusion!
 - Maladies infectieuses, métaboliques, cardio-vasculaires, respiratoires, hématologiques, immunologiques, endocriniennes, ...
 - Troubles du sommeil
 - Troubles du comportement alimentaire
 - Affections psychiatriques

Carfagno DG, et al. Curr Sports Med Rep 2014

Diagnostic

TABLE 1. Diagnosis of OTS—checklist.

Performance - Fatigue	
<p>Is the athlete experiencing the following:</p> <ul style="list-style-type: none"> Unexplainable underperformance Persistent fatigue Increased sense of effort in training Sleep disorders ... <p>Exclusion criteria</p> <p>Are there confounding diseases?</p> <ul style="list-style-type: none"> Anemia Epstein–Barr virus Other infectious diseases Muscle damage (high CK) Lyme disease Endocrinological diseases (diabetes, thyroid, adrenal gland, ...) Major disorders of eating behavior Biological abnormalities (increased erythrocyte sedimentation rate, C-reactive protein, creatinine, or liver enzymes, decreased ferritin, ...) Injury (musculoskeletal system) Cardiological symptoms Adult-onset asthma Allergies ... 	<p>...</p> <p>Are there training errors?</p> <ul style="list-style-type: none"> Training volume increased (>5%) ($\text{h}\cdot\text{wk}^{-1}$, $\text{km}\cdot\text{wk}^{-1}$) Training intensity increased significantly Training monotony present High number of competitions In endurance athletes : decreased performance at “anaerobic” threshold Exposure to environmental stressors (altitude, heat, cold, ...) ... <p>Other confounding factors:</p> <ul style="list-style-type: none"> Psychological signs and symptoms (disturbed POMS, RESTQ-Sport, RPE, ...) Social factors (family, relationships, financial, work, coach, team, ...) Recent or multiple time zone travel ... <p>Exercise test</p> <ul style="list-style-type: none"> Are there baseline values to compare with? (performance, heart rate, hormonal, lactate, ...) Maximal exercise test performance Submaximal or sport-specific test performance Multiple performance tests ...

Meeusen R, et al. *Med Sci Sports Exerc* 2013

Diagnostic – Test de performance

Altération de paramètres physiologiques

- ↓ du time to fatigue
- ↓ VO2max

Altération de paramètres physiologiques CV

- ↓ FC sub max
- ↓ FC max

Biomarqueurs

- ↓ production max de lactate sanguin

Diagnostic – Biomarqueurs

Pas de bon biomarqueur pour le diagnostic de NFOR et OTS

Repos: ↑ CK

A l'exercice: ↓ production max de lactate sanguin à l'exercice

Post-exercice: ⊘ ↑ de la prolactine, l'ACTH, la GH, du cortisol

Diagnostic

Test de performance

Altération de paramètres physiologiques CV

Biomarqueurs (lactate)

Perturbations de l'humeur

Diagnostic – Perturbations de l’humeur

Profile of Mood States (POMS)

- 65 items: fatigue, vitalité, irritabilité, dépression, anxiété/tension, confusion

Daily Analysis of Life Demands for Athletes (DALDA)

Recovery-stress questionnaire for athletes (REST-Q)

- 77 items: stress général, stress émotionnel, stress social, conflits/pression, fatigue, manque d'énergie, plaintes physiques, récupération, stress spécifiques au sport

Training Distress Scale (TDS)

- 7 items: dépression, irritabilité

Questionnaire de dépistage Société Française de Médecine du Sport

Diagnostic – Perturbations de l’humeur

FIGURE 2—Magnitude of changes in POMS mood states from easy to maximal overload training in collegiate varsity swimmers who develop OTS or remain free of symptoms (i.e., “healthy”). (Adapted from Raglin and Morgan [114]).

Meeusen R, et al. Med Sci Sports Exerc 2013

Diagnostic

Test de performance

Altération de paramètres physiologiques CV

Biomarqueurs (lactate)

Perturbations de l'humeur (POMS)

Altérations du métabolisme de base / de la composition corporelle

Atteinte de l'axe hypothalamo-hypophysaire (TOP test)

Titre

FIGURE 1—Relative changes of cortisol, ACTH, PRL and GH responses to the TOP test for normally trained (solid lines), NFO (dashed lines) and OTS (pointed lines) group. Data are presented as percentage increase from both baseline values (SE) of the mean. *Significant main effect of diagnosis $P < 0.05$.

TABLE 1. Sensitivity for detecting NFO vs OTS based on the cutoff values determined by Meeusen et al. (3).

Measure	Cutoff for OTS	OTS ($n = 15$)	NFO ($n = 39$)
Cortisol response to second exercise test	<200% increase*	93% [80%*]	15% [60%*]
ACTH response to second exercise test	<200% increase*	67% [80%*]	74% [80%*]
PRL response to second exercise test	<200% increase*	93% [100%*]	74% [100%*]
GH response to second exercise test	<1000% increase*	47% [40%*]	62% [40%*]

*Cutoff values by Meeusen et al. (3).

Buyse L, et al. *Med Sci Sports Exerc* 2019

Diagnostic

1. Perte de performance sport-spécifique malgré un repos adéquat de 14-21 jours
2. Des signes et symptômes physiques +/- psychologiques compatibles
3. Absence d'autre cause → Diagnostic d'exclusion!
4. Examens paracliniques
 - Test de performance
 - Altération de paramètres physiologiques CV
 - Biomarqueurs (lactate)
 - Troubles de l'humeur (POMS)
 - Altérations du métabolisme de base / de la composition corporelle
 - Atteinte de l'axe hypothalamo-hypophysaire (TOP test)

Traitement

Cas cliniques

H 49ans

Repos:

- Quelques jours off

Éducation:

- Gestion de la charge
- Périodisation
- Stratégies de récupération

F 35ans

Repos

Éducation

Multiples stressseurs:

- Famille, travail
- Alimentation:
 - Shake multivitaminé comme petit déjeuner
 - Repas « légers » sans protéines et peu de féculents
 - Quasi pas de produits laitiers

F 20ans

Repos

Éducation

Multiples stressseurs:

- Trajets
- Période d'examens
- Alimentation pas tout à fait optimale

Mesures préventives

Education

- Planification et périodisation de l'entraînement
- Variations importantes de charges et monotonie = facteurs de risque

Récupération adéquate

- Hydratation, alimentation (CHO, PRO), sommeil, repos
- Monitoring de l'équilibre charge-récupération

Screening médical

- Entraînement, récupération, humeur, stressseurs personnels

Table 5. Preventative measures for nonfunctional overreaching/overtraining syndrome.^{2,44}

Periodization of training
Tapering for competition
Adjust training volume and intensity based on performance and mood
Ensure adequate calories for training load
Ensure adequate hydration
Ensure adequate carbohydrate ingestion during exercise
Ensure adequate sleep
Promoting mental toughness or resilience as buffer
Rest period of greater than 6 hours between exercise bouts
Abstinence of training following infection, heat stroke/stress, periods of high stress
Avoid extreme environmental conditions
Utilize Profile of Mood States (or stress level) and alter training load

Kreher JB, et al. Sports Health 2012

Mesures préventives

Monitoring de certains paramètres

- Charge subjective (méthode Foster)
- FC à l'effort
- Questionnaires psychologiques (POMS) - 1x/mois
- Mesure de performance avec lactatémie - 1x/trimestre
- Métabolisme de base, ou données anthropométriques
- Eventuels biomarqueurs, hormones (?)
- Tests psychomoteurs ralentis (?)

Table 5. Preventative measures for nonfunctional overreaching/overtraining syndrome.^{2,44}

Periodization of training
Tapering for competition
Adjust training volume and intensity based on performance and mood
Ensure adequate calories for training load
Ensure adequate hydration
Ensure adequate carbohydrate ingestion during exercise
Ensure adequate sleep
Promoting mental toughness or resilience as buffer
Rest period of greater than 6 hours between exercise bouts
Abstinence of training following infection, heat stroke/stress, periods of high stress
Avoid extreme environmental conditions
Utilize Profile of Mood States (or stress level) and alter training load

Kreher JB, et al. Sports Health 2012

Figure 4 Triggers, pathophysiology and consequences of overtraining syndrome (OTS).

Figure 5 Predictive model for the identification of overtraining syndrome (OTS). EROS, Endocrine and Metabolic Responses on Overtraining Syndrome; GH, growth hormone; y/o, years old. *Cadegiani FA, et al. BMJ Open Sport Exerc Med 2019*

In athletes that are currently training, particularly during programs or protocols of progressive increasing of training intensity and volume

Fig. 2 Summary of the predictions of Overtraining syndrome (OTS) and its implications

Cadegiani FA, et al. BMC Sports Sci Med Rehabil 2019

Conclusion

Diagnostic d'exclusion

Maladaptation à l'exercice

- Inflammation systémique avec des répercussions neuro-hormonales et immunologiques
- Etat hyporéactif et hypométabolique avec une diminution de la réponse hormonale à l'exercice
→ état anti-anabolique et pro-catabolique

Attentif

- Méthodologie d'entraînement, récupération, stressseurs additionnels, (év. monitoring, screening médical)

Traitement passe par le repos, l'accompagnement, l'éducation, et le traitement des facteurs confondants

Dénominations alternatives?

Underperformance syndrome

Syndrome de déconditionnement paradoxal de l'athlète

Cas cliniques

H 49ans

Amélioration dans semaines suivantes

Optimisation de son régime d'entraînement

F 35ans

A continué à s'entraîner

Repos forcé en raison d'une douleur de hanche

N'a pas pu reprendre ses volumes d'entraînements depuis

F 20ans

Adaptation de l'alimentation

Réduction transitoire du volume d'entraînement

Pas couru le marathon

Résolution quelques semaines après la période d'examens

Pause pendant l'été et augmentation progressive du volume ensuite

Merci pour votre attention

Références utiles

- Bellinger P, et al. Functional Overreaching in Endurance Athletes: A Necessity or Cause for Concern? Sports Medicine, 2020.
- Cadegiani FA, et al. Novel insights of overtraining syndrome discovered from the EROS study. BMJ OpenSport & Exercise Medicine, 2019.
- Cadegiani FA, et al. Novel causes and consequences of overtraining syndrome: the EROS-DISRUPTORS study. BMC Sports Science, Medicine and Rehabilitation, 2019.
- Buyse L, et al. Improving the Diagnosis of Nonfunctional Overreaching and Overtraining Syndrome. Medicine & Science in Sports & Exercise, 2019.
- Cadegiani FA and Kater, et al. Hormonal aspects of overtraining syndrome: a systematic review. Medicine and Rehabilitation, 2017.
- Cardoos N, et al. Overtraining Syndrome. Current Sports Medicine Reports, 2015.
- Carfagno DG, et al. Overtraining Syndrome in the Athlete: Current Clinical Practice. Current Sports Medicine Reports, 2014.
- Meeusen R, et al. Prevention, Diagnosis, and Treatment of the Overtraining Syndrome: Joint Consensus Statement of the European College of Sport Science and the American College of Sports Medicine. Medicine & Science in Sports & Exercise, 2013.
- Birrer D, et al. Prevalence of non-functional overreaching and the overtraining syndrome in Swiss elite athletes. Schweizerische Zeitschrift für Sportmedizin und Sporttraumatologie, 2013.
- Kreher JB, et al. Overtraining Syndrome: A Practical Guide. Sports Health, 2012.